

# ON THE COMPOSITION OF TRANSCENDENTAL ENTIRE AND MEROMORPHIC FUNCTIONS

WALTER BERGWEIFER

ABSTRACT. It is proved that  $f(g) - R$  has infinitely many zeros if  $f$  is a transcendental meromorphic,  $g$  a transcendental entire, and  $R$  a non-constant rational function. The exponent of convergence of the sequence of zeros of  $f(g) - R$  is also estimated.

## 1. INTRODUCTION AND MAIN RESULT

The main result of this paper is the following theorem.

**Theorem .** *Suppose that  $f$  is transcendental meromorphic in the plane, that  $g$  is transcendental entire, and that  $R$  is a non-constant rational function. Then the equation  $f(g(z)) = R(z)$  has infinitely many solutions.*

K. Katajamäki, L. Kinnunen, and I. Laine [7] proved this result under the hypothesis that  $f$  has finite order and  $g$  has finite lower order. In fact, they gave a lower bound for the exponent of convergence of the solutions and also dealt with the case that  $R$  is transcendental but of smaller growth than  $g$ . While our method does not seem to be suitable to handle transcendental functions  $R$ , it does give a bound for the exponent of convergence, compare §3. The results of K. Katajamäki, L. Kinnunen, and I. Laine [7] generalized their results of [6] where they assumed that  $f$  is entire as well as the result of [3] where the above theorem was proved under the additional hypothesis that  $f(g)$  has finite order.

The above theorem is known in the case that  $R(z) = z$  [2], as well as in the case that  $f$  is entire and  $R$  is a polynomial [1]. These results confirmed a conjecture of F. Gross [5]. The method used in [2] is based on the observation that  $f(g)$  has infinitely many fixpoints if and only if  $g(f)$  does. The underlying idea in the present paper is to consider the solutions of  $f(g(z)) = R(z)$  as fixpoints of  $R^{-1}(f(g(z)))$  and to proceed similarly as in [2]. This requires some modifications of the argument, however, because  $R^{-1}$  is, in general, not a single-valued function.

We note that the method of [1] extends to the case that  $R$  is rational. It does not seem to extend, however, to the case that  $f$  is meromorphic. On the other hand, the case that  $f$  has only one pole can be handled by this method with only minor modifications.

## 2. PROOF OF THE THEOREM

We may assume that  $R(\infty) = \infty$  and in fact that  $R(z) \sim z^n$  for some  $n \in \mathbb{N}$  as  $z \rightarrow \infty$  because otherwise we consider  $L(f)$  and  $L(R)$  instead of  $f$  and  $R$  for a

---

1991 *Mathematics Subject Classification.* 30D05, 30D35.

*Key words and phrases.* Meromorphic function, entire function, composition, factorization, fixpoint.

suitable linear transformation  $L$ . In view of the remarks made at the end of the introduction, we may also assume that  $f$  has at least two poles  $z_1$  and  $z_2$ . By  $p_j$  we denote the order of  $z_j$ .

By [2, Lemma 1] there exist functions  $h_j$  analytic in a neighborhood of 0 such that  $h_j(0) = 0$  and  $f(h_j(z) + z_j) = z^{-p_j}$  for  $j = 1, 2$ . Similarly as in [2] we define  $k_1(z) = h_1(z^{p_2 n}) + z_1$  and  $k_2(z) = h_2(z^{p_1 n}) + z_2$  so that  $f(k_1(z)) = f(k_2(z)) = z^{-p_1 p_2 n}$ .

Next we note that  $R(z) = S(z)^n$  for some function  $S$  which is univalent in a neighborhood of  $\infty$  and satisfies  $S(z) \sim z$  as  $z \rightarrow \infty$ . We denote the inverse function of  $S$  by  $T$ . Then  $T(z) \sim z$  as  $z \rightarrow \infty$  and  $R(T(z)) = z^n$ . Finally, following [2], we define  $u(z) = g(T(z^{-p_1 p_2}))$  and

$$v(z) = \frac{u(z) - k_1(z)}{u(z) - k_2(z)}.$$

Then 0 is an essential singularity of  $u$  and hence  $v$ . Because  $k_1(0) = z_1 \neq z_2 = k_2(0)$  we have  $v(z) \neq 1$  for sufficiently small  $z$ . Hence Picard's theorem implies that  $v$  takes one of the values 0 and  $\infty$  in any punctured neighborhood of 0. Without loss of generality we may assume that this holds for the value 0, that is, there exists a sequence  $\zeta_j$  tending to 0 such that  $v(\zeta_j) = 0$ . It follows that  $u(\zeta_j) = k_1(\zeta_j)$  and hence that

$$f(g(T(\zeta_j^{-p_1 p_2}))) = f(u(\zeta_j)) = f(k_1(\zeta_j)) = \zeta_j^{-p_1 p_2 n} = R(T(\zeta_j^{-p_1 p_2})),$$

that is,  $f(g(\omega_j)) = R(\omega_j)$  for  $\omega_j = T(\zeta_j^{-p_1 p_2})$ . The conclusion follows since  $\zeta_j \rightarrow 0$  implies that  $\omega_j \rightarrow \infty$ .

### 3. A QUANTITATIVE VERSION OF THE MAIN RESULT

Denote by  $\rho(f)$  and  $\lambda(f)$  the order and the lower order of  $f$  and by  $\sigma$  the exponent of convergence of the zeros of  $f(g(z)) - R(z)$ . K. Katajamäki, L. Kinnunen, and I. Laine [7] have shown that  $\sigma \geq \lambda(g)$ , provided  $\lambda(g) < \infty$  and  $\rho(f) < \infty$ .

We shall show that the slightly stronger inequality  $\sigma \geq \rho(g)$  can be obtained without growth restrictions on  $f$  or  $g$  if we make the assumptions of §2:

*If  $f$  has at least two poles and  $R(\infty) = \infty$ , then  $\sigma \geq \rho(g)$ .*

To prove this result, we proceed as in §2. Instead of Picard's theorem, however, we use Nevanlinna theory for functions meromorphic in the neighborhood of an essential singularity, compare e. g. [4, §78ff.]. By  $\rho(u)$  and  $\rho(v)$  we denote the orders of  $u$  and  $v$  at 0. It is easily seen from the definitions of  $u$  and  $v$  that  $\rho(v) = \rho(u) = p_1 p_2 \rho(g)$ . By Borel's theorem [4, p. 354], we may assume that the exponent of convergence of  $\zeta_j$  at 0 is equal to  $\rho(v)$  so that  $\sum_{j=1}^{\infty} |\zeta_j|^{-\mu}$  diverges for all  $\mu < \rho(v)$ . Because  $\omega_j = T(\zeta_j^{-p_1 p_2}) \sim \zeta_j^{-p_1 p_2}$  as  $j \rightarrow \infty$  this implies that  $\sum_{j=1}^{\infty} |\omega_j|^{-\mu}$  diverges for all  $\mu < \rho(g)$ , that is, the exponent of convergence of the zeros of  $f(g) - R$  is at least  $\rho(g)$ .

### 4. REMARKS

The estimate  $\sigma \geq \rho(g)$  is probably far from being best possible. It seems likely to me that the counting function of the zeros of  $f(g) - R$  and the Nevanlinna characteristic of  $f(g)$  are always of the same order of magnitude. Possibly the Nevanlinna deficiency  $\delta(0, f(g) - R)$  is always equal to 0 if  $f$ ,  $g$ , and  $R$  are as in the statement of our main theorem. We remark that J. K. Langley [8] proved that if  $f$

and  $g$  are entire transcendental and if  $f(g)$  is of finite order, then  $\delta(0, f(g) - R) < 1$  for any non-constant rational function  $R$ . For further results concerning the number of zeros of  $f(g) - R$  for entire  $f$  we refer to C.-C. Yang and J.-H. Zheng [10].

Finally we mention that the conclusion of our main theorem remains valid if  $f$  is a rational function of degree at least 2, see K. Katajamäki, L. Kinnunen, and I. Laine [7] or G. S. Prokopovich [9]. This follows also from our proof if  $f$  has at least two poles.

## REFERENCES

- [1] W. Bergweiler, Proof of a conjecture of Gross concerning fix-points, *Math. Z.* 204 (1990), 381-390.
- [2] W. Bergweiler, On the existence of fixpoints of composite meromorphic functions, *Proc. Amer. Math. Soc.* 114 (1992), 879-880.
- [3] W. Bergweiler and C.-C. Yang, On the value distribution of composite meromorphic functions, *Bull. London Math. Soc.* 25 (1993), 357-361.
- [4] A. Dinghas, *Vorlesungen über Funktionentheorie*, Springer, Berlin, Göttingen, Heidelberg 1961.
- [5] F. Gross, *Factorization of meromorphic functions*, U.S. Government Printing Office, Washington, DC, 1972.
- [6] K. Katajamäki, L. Kinnunen, and I. Laine, On the value distribution of composite entire functions, *Complex Variables Theory Appl.* 20 (1992), 63-69.
- [7] K. Katajamäki, L. Kinnunen, and I. Laine, On the value distribution of some composite meromorphic functions, *Bull. London Math. Soc.* 25 (1993), to appear.
- [8] J. K. Langley, On the fixpoints of composite entire functions of finite order, *Proc. Royal Soc. Edinburgh*, to appear.
- [9] G. S. Prokopovich, Fix-points of meromorphic functions, *Ukrain. Math. J.* 25 (1973), 198-208; translation from *Ukrain. Mat. Zh.* 25 (1973), 248-260.
- [10] C.-C. Yang and J.-H. Zheng, Further results on fixpoints and zeros of entire functions, *Trans. Amer. Math. Soc.*, to appear.

LEHRSTUHL II FÜR MATHEMATIK, RWTH AACHEN, D-52056 AACHEN, GERMANY  
*E-mail address:* bergw@math2.rwth-aachen.de